

Mechanisms and mediators of hypertension induced by erythropoietin and related molecules

Rajiv Agarwal

Division of Nephrology, Department of Medicine, Indiana University School of Medicine and Richard L. Roudebush Veterans Administration Medical Center, Indianapolis, IN, USA

Correspondence and offprint requests to: Rajiv Agarwal; E-mail: ragarwal@iu.edu

ABSTRACT

Hypertension is a common but frequently overlooked adverse effect of erythropoietin (EPO) therapy. Underreporting of hypertension with EPO is likely due to either more aggressively managing hypertension through the prescription of antihypertensive drugs or closer attention to dry weight. The purpose and focus of this review is to critically evaluate the mechanisms of EPO-induced hypertension. Preclinical data are considered first, followed by clinical data where available. Mediated by a variety of molecules, there is an imbalance in the vascular tone favoring net vasoconstriction that mediates EPO-induced hypertension. Animal studies show the primary importance of chronic kidney disease in the genesis of EPO-induced hypertension. Preclinical studies show deranged regulation of the nitric oxide, endothelins and prostanoids and the sympathoadrenal and renin-angiotensin pathways as causes of EPO-induced hypertension. Human studies suggest that EPO administration is also associated with increased responsiveness to catecholamines and angiotensin II on vascular tissue; in addition, hypoxia-induced vasodilation may be impaired in those with EPO-induced hypertension. There is little evidence for EPO as a direct vasoconstrictor or its effect on blood viscosity as a mechanism of EPO-induced hypertension. EPO-induced hypertension, at least in part, appears to be independent of an increase in hemoglobin, because experiments show that hemoglobin may be increased by EPO without an increase in blood pressure (BP) by simply treating the animals with EPO-binding protein and that treatment with EPO in the setting of iron deficiency may not increase hemoglobin but may still increase BP. However, experimental data are not consistent across studies and better mechanistic designs are needed, especially in patients with chronic kidney disease, to dissect the precise mechanism of EPO-induced hypertension. Animal studies suggest that hypoxia-inducible factor stabilizers may induce hypertension by provoking calcification and augmenting chronic intermittent hypoxia as occurs in sleep apnea. Others show that there may be an antihypertensive effect via kidney repair.

Whether these drugs will alter the risk of hypertension compared with EPO remains to be seen.

Keywords: anemia, erythropoietin, HIF stabilizers, hypertension, nitric oxide

Hypertension is a common but frequently overlooked adverse effect of erythropoietin (EPO) therapy [1–6]. Underreporting of hypertension with EPO is likely due to either more aggressively managing hypertension through the prescription of antihypertensive drugs, closer attention to dry weight [7, 8] or lack of ambulatory blood pressure (BP) measurements [9]. The purpose and focus of this review is to critically evaluate the mechanisms of EPO-induced hypertension. Preclinical data are considered first, followed by clinical data where available.

IMPORTANCE OF CHRONIC KIDNEY DISEASE (CKD) IN EPO-INDUCED HYPERTENSION

The first study to show that CKD was an important determinant for EPO-induced hypertension was reported from France in 1995 [10]. Male Wistar rats underwent subtotal nephrectomy at 6 weeks of age, were allowed to recover for 6 weeks and were then treated for 3 weeks with either vehicle or EPO. Time controls were sham-operated animals that did not undergo nephrectomy and were treated with either vehicle or EPO. As expected, subtotal nephrectomy in animals increased plasma creatinine and systolic BP and induced a decrease in hematocrit. Treatment with EPO provoked a significant increase in hematocrit regardless of nephrectomy. EPO treatment increased systolic BP significantly in the nephrectomized group by 37.5 ± 11 mmHg. There was no significant increase in systolic BP in the nonnephrectomized EPO-treated or vehicle-treated rats. Furthermore, no relationship was seen between the change in hematocrit and the change in systolic BP [10]. Using a very similar design as above, a 1997 study from Quebec confirmed the

central importance of reduced renal mass in the development of hypertension [11]. EPO had no effect on BP in the control rats despite an increase in hematocrit in rats without nephrectomy, yet had a large effect in those with a nephrectomy.

The subtotal nephrectomy model provides a consistent observation for both studies in that by itself—over the time course studied—it did not provoke hypertension. Furthermore, there was no increase in BP in the EPO-treated sham-operated animals. It required both subtotal nephrectomy and EPO to provoke hypertension. Whether nephrectomy activated the renin–angiotensin system, caused volume excess or caused endothelial dysfunction to induce hypertension was not examined in these studies, but these possibilities were subsequently explored in other models and will be discussed further below.

Human observations are consistent with animal data. Although EPO is abused by some athletes to enhance performance, hypertension or hypertensive crises are not reported. Similarly, the incidence of hypertension in zidovudine-treated human immunodeficiency virus–positive anemic patients or among those undergoing cancer chemotherapy is low. However, EPO-induced hypertension is very common among those with CKD, including those on dialysis [6].

HEMOPOIETIC VERSUS HYPERTENSIONOGENIC EFFECTS OF EPO

Whether BP tracks with hemoglobin or not remains controversial. In a 1988 report, among anemic rats with reduced renal mass, preventing anemia of renal failure with EPO aggravated systemic hypertension [12]. In contrast, a low-iron diet that maintained hemoglobin at a lower level prevented the development of hypertension, hyperfiltration, glomerulosclerosis and proteinuria [12]. In more recent animal experiments, hypertension did not track with an increase in hemoglobin [13]. If EPO is administered to anemic animals with chronic renal failure, but hemoglobin is kept stable by feeding an iron-deficient diet, hypertension still occurs. In blood vessels harvested from these animals treated with EPO, vasodilatory responses to nitric oxide (NO) donors were impaired but response to several vasoconstrictors was normal. Among patients on long-term hemodialysis, treatment with iron to increase hemoglobin was not associated with parallel increases in BP [14].

Human data are similarly inconclusive. Some studies show that although an increase in hemoglobin is dose dependent, an increase in BP is not [1, 15]. On the other hand, during the first 5 weeks of administration of EPO the change in hemoglobin concentration was directly related to an increase in diastolic BP ($r = 0.42$, $P < 0.001$) [3].

To understand the hypertensionogenic effects of EPO, consideration of the biology of the EPO receptor expressed in the nonhemopoietic tissues is needed.

EPO RECEPTOR AND ITS RELEVANCE TO EPO-INDUCED HYPERTENSION

Mutation of EPO receptor is lethal *in utero* [16]. However, EPO receptor null mice can survive simply by expressing the EPO receptor in the hemopoietic tissue only [16]. Despite the lack of

the EPO receptor in the endothelium, heart and brain, these mice, somewhat surprisingly, develop normally. Using genetic engineering to rescue EPO receptor null mice, two groups of mice were created with varying amounts of EPO receptor expressed in the hemopoietic tissue [16]. In one group, the expression of EPO receptor was 40% of normal and in the second group was 120% of normal. As expected, the circulating EPO level was higher when the EPO receptor was lower. However, contrary to expectations, in response to induced anemia the time to peak plasma EPO concentrations was delayed in both groups of mice. This suggests that the extra-hemopoietic EPO receptor may play an important role in the regulation of plasma EPO concentration.

There is substantial evidence that blood vessels express the EPO receptor. In a 1994 report, mRNA for the EPO receptor was identified in human umbilical vein endothelial cells [17]. These endothelial cells lacked erythroid precursors because they lacked the α -globin and γ -globin chains associated with hemoglobin. Furthermore, antibodies to the extracellular portion of the EPO receptor stained the vascular endothelium of the placenta and the umbilical cord. Cultured endothelial cells proliferate in response to EPO [18, 19]. The ability to proliferate is seen at concentrations of EPO as low as 1 U/mL [19], which is often achieved with intravenous administration of the drug [20]. The ability of human umbilical vein endothelial cells to migrate is also induced by EPO [19].

A series of studies performed by investigators in Japan among mice expressing EPO receptor in only the hemopoietic tissue demonstrate the importance of EPO receptor expression outside hemopoietic tissue [16] such as heart, lungs and the limbs [21]. Experiments in these mice lacking the EPO receptor outside hemopoietic tissue display several cardiac, pulmonary and vascular anomalies as follows: (i) Following coronary ligation, compared with wild-type mice, there was greater infarct size, due in part to accelerated cardiomyocyte apoptosis [22]. (ii) When pressure overload was induced by aortic banding, there was greater cardiac dilatation and impairment in cardiac contractility, due in part to reduced vascular endothelial growth factor expression and reduced myocardial capillary density [23]. In response to hypoxia, there was an increased risk of the development of pulmonary hypertension due to reduced mobilization of the endothelium progenitor cells [24]. (iv) In response to hind-limb ischemia, there was abrogation of the expression of vascular endothelial growth factor and its receptor, impairment in the recruitment of endothelial progenitor cells and bone marrow–derived proangiogenic cells and thereby reduced neovascularization [25].

Some data suggest that the effects on hemopoiesis and vasoconstriction may be mediated by different epitopes on the EPO molecule [26]. Rodents treated with EPO, when also treated with an EPO binding protein, had an increase in hemoglobin but their BP did not change. Furthermore, treatment of the above animals with the EPO binding protein and an antibody to this protein increased hemoglobin but did not change BP. These data suggest that the BP increasing effect of EPO and the hemopoietic effect of the EPO molecule may be mediated by different parts of the same molecule.

EPO can also be engineered by targeting it more specifically to the hemopoietic EPO receptor and reducing its ability to

bind to nonhemopoietic tissues [27]. A synthetic EPO molecule has been created that interacts minimally with the EPO receptor [27]. To selectively target EPO to the hemopoietic tissue, it was tethered to an antibody that specifically binds glycophorin A, which is highly expressed on red blood cells [27]. Compared with wild-type mice, when this new molecule was given to mice carrying the human glycophorin A gene, the half-life of engineered EPO was prolonged, reticulocyte response was augmented and platelet effects were minimized [27]. The authors speculate that EPO receptor expression on maturing megakaryocytes may create an off-target prothrombotic state. In their study, since the total platelet count and reticulated platelets were both reduced in comparison to darbepoietin, the new molecule may reduce the risk of thrombotic side effects. The promise of an engineered EPO with less BP raising effect has yet to be realized.

MOLECULAR MECHANISMS OF EPO-INDUCED HYPERTENSION

The mechanisms of EPO-induced hypertension are incompletely understood. Figure 1 summarizes putative mediators and their effects on BP are discussed further.

Evidence against direct effect of EPO on vascular smooth muscle or endothelium

Among untreated patients with essential hypertension, serum EPO concentration correlates with both systemic

vascular resistance and 24-h ambulatory BP [28]. This has led to speculation that EPO may have direct vasoconstrictive effects. In order to interpret the results of cell culture studies, it is important to compare the plasma concentrations of EPO achieved after a bolus dose in humans with that used in preclinical studies. A study describing the pharmacokinetics of EPO in hemodialysis patients notes that the peak concentration of EPO when given as an intravenous bolus injection was 0.768 U/mL with 50 U/kg EPO and 2.434 U/mL with 150 U/kg EPO [20].

In 1991, a group of investigators from Muenster, Germany demonstrated that resistance arterioles from the kidneys and intestine could be made to vasoconstrict when exposed to EPO in doses ranging from 10 to 200 U/mL [29]. In 1993 the same group demonstrated that intracellular calcium is increased in cultured vascular smooth muscle cells in a dose-dependent manner upon incubation with EPO [30]. The vascular smooth muscle cells were incubated with 100 U/mL or 250 U/mL EPO. It should be noted that these concentrations of EPO are ~100–200-fold higher than the peak concentrations achieved with intravenous EPO administration [20]. The physiological and clinical relevance of these observations is therefore unclear.

In humans, EPO by itself does not appear to have any effect on vasoconstriction. In a double-blind cross-over study in nine hemodialysis patients, Hon *et al.* [31] administered either EPO or saline intravenously. BP was measured every 5 min for 60 min following treatment. Between treatments, no differences




FIGURE 1: Putative mechanisms of EPO-induced hypertension through its receptors on the endothelial cells, especially in the setting of CKD, can trigger endothelial dysfunction and vasoconstriction. The NO pathway is severely deranged in response to EPO; ADMA concentrations increase and cGMP release with NO donors is impaired. Inhibitors of NOS severely elevate BP and provoke mortality in rodents with excess EPO expression. NO can also trigger ET-1 release, which itself is a potent vasoconstrictor. Prostanoids are downstream to endothelin and there is a net imbalance favoring vasoconstriction. The effects on the renin–angiotensin system are complex, but in preliminary studies vasoconstriction to angiotensin II is enhanced when patients are given EPO. Similarly, compared with patients not on EPO, vasoconstriction to catecholamines is enhanced when patients are treated with EPO. Additional pathways such as volume excess and other mechanisms discussed in the text are not depicted in the figure.

were seen [31]. A 2004 study in 41 dialysis patients reported an increase in mean arterial pressure from 103 to 105 mmHg after a single injection [32]. Although the change was only 2 mmHg, it was reported as significant; without a blinded control group, the likelihood of false discovery is high [32]. A 2013 study used subcutaneous arteries isolated from a gluteal biopsy from 17 patients with Stage 4 CKD [33]. The response to endothelial-dependent vasodilatation was tested *ex vivo* by evaluating acetylcholine-induced vasodilatation. These blood vessels were exposed to varying EPO concentrations at 0, 1, 10 and 20 IU/mL [33]. The difference in endothelial function was minimal when comparing 0 and 1 IU/mL but stark at higher concentrations. However, such higher concentrations are supra-pharmacologic and it is unclear if they have relevance to EPO biology. Furthermore, the effects of EPO may be mediated via endothelin, because using an endothelin type A receptor antagonist abrogated the effects of EPO [33].

Endothelin-1

Rats with reduced renal mass have increased endothelin-1 (ET-1) expression in the aorta, mesenteric artery and renal cortex [34, 35]. These rats have increased 24-h urinary excretion of ET-1 [36]. EPO induces release of ET-1 from endothelial cells in culture [18, 37]. In contrast to the cell culture experiments, EPO administered to 5/6 nephrectomized rats does not increase the circulating concentration of ET-1 despite an increase in BP [38, 39]. This may be because the expression of ET-1 provoked by EPO is organ specific. EPO-treated rats with reduced renal mass have no increase in ET-1 in the mesenteric artery or renal cortex, yet the aortic content of ET-1 is increased [39, 40]. The increase in aortic ET-1 was further augmented by the NO synthase (NOS) inhibitor L-arginine analog N^G-nitro-L-arginine methyl ester (L-NAME). This suggests that NO has an important effect on abrogating the expression of ET-1 in large vessels and is an upstream process [40]. Abrogation of oxidative stress in EPO-treated nephrectomized rats by tempol also reduces tissue levels of ET-1, hypertension and renal injury [41]. This suggests that oxidative stress is also an upstream process to ET-1-generation in the kidney. However, as discussed below, prostanoids may be a downstream mediator of ET-1 induced vasoconstriction. Mice specifically expressing ET-1 in the endothelium, when treated with EPO even without nephrectomy, have an increase in BP, impairment of endothelial function, resistance artery remodeling and aortic inflammation and oxidative stress [42]. These adverse effects can be blocked with exercise over 8 weeks.

Consideration of the tissue-specific expression of endothelin receptors is needed to appreciate the endothelin biology. Rats with reduced renal mass have reduced expression of the vasodilatory endothelin receptor type B (ET_B) in the aorta, mesenteric artery and renal cortex [34, 35]. Furthermore, in rats with reduced renal mass, EPO increases the vascular and renal cortex expression of ET_B receptor mRNA [43]. This parallels the increase in ET_B receptor density in the endothelium of rats [43]. Compared with wild-type mice, mice heterozygous for the ET_B receptor have a greater increase in BP with EPO [43]; mice heterozygous for the ET_A receptor have BP responses that are similar to wild-type mice. In rats with subtotal nephrectomy and

EPO-induced hypertension, treatment with a selective ET_A receptor antagonist is more effective than placebo in abrogating the increase in BP [44]. In contrast, the nonselective ET_A/ET_B receptor antagonist, bosentan had no effect in preventing EPO-induced hypertension.

In patients with CKD treated chronically with EPO, ET-1 concentration is increased [45–48]. In a cross-sectional study of 44 end-stage renal disease patients [24 hemodialysis, 20 continuous ambulatory peritoneal dialysis (CAPD)], of which half were on EPO therapy, plasma ET-1 concentration directly correlated with systolic BP [49]. In EPO-treated patients on CAPD, plasma ET-1 concentrations correlated with mean BP [50]. This correlation is not seen in those not treated with EPO [50]. However, the acute administration of EPO does not increase ET-1 concentration [31]. Vasoconstriction induced by infusion of ET-1 is similar in patients with CKD before and after treatment with EPO, thus the vasoconstriction dose–response relationship to ET-1 is not altered with EPO-treatment [51].

In summary, treatment of anemia of CKD with EPO uncovers the endothelial dysfunction through increased production of vascular ET-1. Reduced expression of the vasodilatory ET_B receptor due to genetic heterogeneity or cross talk by other endothelium-derived vasoactive autacoids such as NO, prostanoids and angiotensin II may play an important role in the genesis of EPO-induced hypertension [35].

Prostanoids

Prostanoids play an important role in maintaining vascular tone and renal sodium handling. Thromboxane A₂ (TXA₂) and prostaglandin F_{2α} (PGF_{2α}) have vasoconstrictor activity, whereas PGI₂ is vasodilatory. TXB₂ is a stable metabolite of TXA₂, whereas 6-keto-PGF_{1α} is a stable metabolite of PGI₂ [37]. In a 2005 study, rats with reduced renal mass who became anemic and hypertensive as expected also had an increase in vascular and renal concentration of TXA₂ and PGI₂ [52]. In these animals, treatment with EPO further aggravated hypertension and stimulated vascular and renal TXA₂ and PGI₂ [52]. Pretreatment with aspirin blocks both the synthesis of TXA₂ and PGI₂ and therefore had no net effect on EPO-induced hypertension [52].

In contrast to systemic exposure of EPO, rabbit aortic rings incubated with EPO show an increase in TXA₂ but a reduction in PGI₂ [18, 37]. In human renal artery *in vitro* as well as rabbit aortic rings, incubation with EPO leads to an increase in the synthesis of constrictor prostanoids such as TXB₂ and PGF_{2α} but a reduction in the vasodilatory prostaglandin PGI₂ [53]. However, the concentration of EPO used to incubate these rings was far beyond the physiological range. TXB₂ was increased and 6-keto-PGF_{1α} unchanged in the aorta of rats with reduced renal mass treated with EPO [54]. An antagonist of thromboxane, ridogrel, abrogated hypertension in EPO-treated animals but did not alter the plasma concentration of ET-1 [54]. An antagonist of ET-1, ABT-627, was even more effective than ridogrel in treating hypertension; moreover, it also reduced the concentration of TXB₂ in the aorta [54]. Thus prostanoids may serve as a downstream mediator of vasoconstriction in response to ET-1 activation. The imbalance in vasoconstrictor to vasodilatory prostanoids may lead to a net increase in vascular

resistance and therefore hypertension. The use of antiplatelet therapy has been postulated to prevent the development of hypertension among patients treated with EPO [55]; the mechanism of the antihypertensive effect remains unclear.

NO

NO is a potent endothelium-derived vasodilator and plays an important role in the genesis of hypertension in CKD. NO has important effects on the endothelium, blood vessels, kidneys and BP. Each of the effects are discussed below.

Endothelium. In uremia, NO synthesis in the endothelium is blunted. Suprapharmacologic EPO exposure to endothelial cells in culture media results in a reduction in endothelial NOS [56] and high concentrations (such as 10–200 U EPO/mL) provoke an increase in the NOS inhibitor asymmetric dimethylarginine (ADMA) in a dose-dependent manner; this is accompanied by reduced activity of dimethylarginine dimethylaminohydrolase (DDAH) [57]. DDAH is an enzyme that breaks down ADMA, generating citrulline and dimethylamine, and is the major mechanism of ADMA removal [58]. *In vivo* studies reveal that compared with saline-treated controls, Balb/c mice with intact kidneys injected with more reasonable exposure to EPO (such as 30 U every other day for 10 weeks) show ADMA concentrations that were 46% higher [59]. In comparison, symmetric dimethyl arginine (SDMA) concentrations are unchanged. Of note, there is no mechanism of direct synthesis of ADMA; free ADMA is generated by proteolysis of methylated proteins [58]. Free ADMA can inhibit endothelial cell NOS and cause endothelial dysfunction and has been linked to poor outcomes in CKD.

Blood vessels. In rabbit, aortic rings harvested from animals with intact kidneys, acetylcholine-induced vasodilatation, which is dependent upon an intact endothelium, was not blunted when rings were incubated with EPO for 30 min [53]. In sharp contrast, among rats with reduced renal mass, the release of cyclic guanosine monophosphate (cGMP) by nitrate donors of aortic rings is augmented. However, this augmentation is blunted when rats were treated with EPO [60]. The endothelial and inducible NOS protein mass is reduced in these rats regardless of EPO treatment, but when treated with felodipine, the protein mass in the aorta is restored. Thus the presence or absence of renal failure has an important effect on the role of EPO in blood vessels; calcium-channel blockade may have at least partially restored the abnormalities.

Kidney effects. Rats with intact renal mass upon treatment with EPO display an increase in renal NO; this is noted by an increase in urinary cGMP [61] and urinary nitrates [62]. Rats with reduced renal mass have a reduction in the urinary nitrate excretion rate that is further reduced with EPO treatment [63]. The endothelial and inducible NOS protein mass is reduced in rats with subtotal nephrectomy regardless of EPO treatment, but when treated with felodipine the protein mass is restored [63]. When mice with intact renal mass are given EPO, although ADMA concentrations in plasma are increased, the urinary excretion of NO metabolites is also increased [59].

Similar findings are seen in rats with intact kidneys treated with EPO [64]. In fact, mice treated with EPO for 10 weeks have an increased kidney expression of NOS1 and NOS2 despite the EPO-induced increase in ADMA [59]. Similarly, rats with intact kidneys made hypertensive with EPO have increased urinary excretion of nitrites and nitrates [65]. The renal vasodilatory response to acetylcholine (endothelium dependent) and sodium nitroprusside (endothelium independent) are both similar to vehicle-treated rats [65]. Thus, reduced renal mass is necessary to uncover the effects of EPO.

BP effects. Inhibitors of NOS can exacerbate hypertension in EPO-treated animals even when the renal mass is not reduced [61]. Among animals treated with EPO, exposure to L-NAME, an inhibitor of NOS, results in hypertension [40], the magnitude of which is greater when renal mass is reduced [40]. In rats in whom CKD is induced by ligating renal arteries, the restoration of endothelial NOS by gene delivery improves NO release and prevents the development of hypertension [66]. In rats with reduced renal mass and EPO-induced hypertension, the treatment of hypertension with felodipine partially restores the abnormalities in NO metabolism [63].

The critical importance of NO in EPO-induced hypertension has been studied in a transgenic mouse model [67]. Transgenic mice overexpressing human EPO were generated. Despite hematocrit levels of 80%, adult transgenic mice did not develop hypertension or thromboembolism because of a compensatory increase in endothelial NOS levels, NO-mediated endothelium-dependent relaxation and circulating and vascular tissue NO levels. Administration of L-NAME led to vasoconstriction, an increase in vascular resistance, hypertension and death of transgenic mice; the wild-type siblings developed hypertension but did not show increased mortality.

A translational research study elucidated the mechanism of endothelial dysfunction in 56 patients on hemodialysis treated with EPO [68]. Endothelial progenitor cells, which the investigators state reflect endothelial cell function, were isolated from these patients and mRNA levels for both the full EPO receptor and a spliced, truncated EPO receptor were measured. The authors note that activation of the full EPO receptor triggers a signaling cascade that ultimately terminates in cGMP and NO production and subsequent vasodilation. However, it was observed that in patients with EPO-induced hypertension there was a positive correlation with the spliced, truncated variant of the receptor. This led to the conclusion that the truncated variant of the receptor serves as a dominant negative regulator of the cGMP/NO cascade, thus blunting vasodilation and possible downstream hypertension.

From the above it appears that the effects on BP due to EPO exposure may be mediated by the generation of endothelial NO. At least there is some evidence for a reduction in NO generation and blunting of the NO effect on both the kidneys and vasculature. Transgenic models demonstrate that endothelial NO appears to be critical in maintaining normotension, preventing cardiovascular dysfunction and survival *in vivo* in the setting of EPO use.

Catecholamines

In rabbit, aortic rings incubated with very high EPO concentrations result in increased vasoconstriction when exposed to norepinephrine [37, 53]. Norepinephrine concentrations were increased following 12 weeks of EPO treatment in one study [69] but decreased in another study of Japanese hemodialysis patients [70]. Vasoconstriction induced by infusion of norepinephrine was increased in patients with CKD after treatment with EPO, thus the vascular sensitivity to norepinephrine was increased in human studies [51, 71]. Furthermore, white blood cell α -2 receptor density was reduced following EPO treatment [45, 72]. In summary, these data suggest both an elevation of catecholamines as well as enhanced sensitivity to catecholamines on the blood vessels as a mechanism of EPO-induced hypertension.

Renin–angiotensin–aldosterone system (RAAS)

Eggena *et al.* [73] evaluated the effects on the RAAS at the molecular level in Wistar rats upon treatment with EPO. In the kidney, both renin mRNA and angiotensinogen mRNA were increased by EPO. In the heart, no alterations in mRNAs were seen. In the aorta, angiotensinogen-mRNA but not renin mRNA was elevated. In both the aorta and kidney, a significant correlation was observed between angiotensinogen mRNA and BP. In cultured rat vascular smooth muscle cells, Barrett *et al.* [74] show that EPO induces an increase in both types of angiotensin II receptors, even in the presence of enalapril or losartan. Among hemodialysis patients, infusion of angiotensin II leads to excess vasoconstriction, suggesting that angiotensin II sensitivity is increased with EPO treatment [51]. These data suggest that the vasoconstrictive potential is enhanced with EPO treatment.

In isolated perfused rodent kidneys, EPO treatment results in RAAS-stimulated sodium retention [75]. This is in sharp contrast to normal human volunteers where EPO treatment caused a reduction in plasma volume, plasma renin activity and aldosterone [76]. However, in rats with subtotal nephrectomy treated with EPO, in terms of treatment there is no unique effect on BP lowering of RAAS blockade. This is because the reduction in systolic BP was similar when rats were treated with traditional triple therapy (reserpine, hydralazine and hydrochlorothiazide) or with the RAAS blockers captopril or losartan [36].

Despite an increase in mean arterial pressure in 12 Japanese hemodialysis patients treated with EPO, plasma renin activity was found to be reduced [70]. Among dialysis patients treated with EPO, a close relationship is seen between exchangeable sodium, an increase in plasma aldosterone and an increase in BP [77]. As discussed above, attention to dry weight can abrogate EPO-induced hypertension among dialysis patients.

The above data—both in animals and humans—do not exclude the possibility of EPO causing hypertension by inducing volume excess or in the setting of volume excess. For example, the rodent subtotal nephrectomy models are associated with volume overload, which may be a prerequisite for developing hypertension. EPO, as noted above, can provoke sodium

retention as well. However, the quality of the data do not allow deducing a cause-and-effect relationship.

Blood viscosity

Blood viscosity increases in parallel with blood hematocrit and has been cited as a mechanism of EPO-induced hypertension. However, not all patients who have correction of anemia get hypertensive. Thus the change in blood viscosity by itself appears to be insufficient to account for EPO-induced hypertension.

Vascular sensing of hypoxia

Breathing 60% oxygen for 10–12 min leads to an increase in forearm vascular resistance paralleled by a reduction in forearm blood flow, but not in all patients [78]. This phenomenon is only observed in those patients in whom BP increases with EPO treatment [78]. Thus the vascular response to hypoxia—and its reversal with correction of anemia—may be a fundamental mechanism of the genesis of hypertension induced by EPO.

HYPOXIA-INDUCIBLE FACTORS: PROLYL HYDROXYLASE INHIBITORS

EPO gene expression is provoked by hypoxia-inducible transcription factors (HIFs) [79]. HIF- α subunits are oxygen labile and, under normal oxygen tension, oxidized by prolyl hydroxylases. Inhibitors of prolyl hydroxylases can stabilize HIFs, simulate hypoxia and promote erythropoiesis [79]. HIF stabilizers not only stimulate EPO, but also many other genes responsible for angiogenesis, tumor growth, cell proliferation and metabolism. HIF stabilizers may aggravate hypertension by several mechanisms. For example, chronic intermittent hypoxia through HIF signaling in the carotid artery is thought to provoke systemic hypertension [79]. A 2016 study in *in vitro* and *in vivo* rodent models shows that hypoxia induces inorganic phosphorus-induced vascular smooth muscle calcification [80]. In this rodent model, roxadustat, an oral HIF prolyl hydroxylase inhibitor, enhanced vascular calcification [80]. The downstream effect of long-term use may therefore be hypertension. Vascular calcification is common in CKD and contributes to arterial stiffness. Increased arterial stiffness is strongly associated with elevated interdialytic ambulatory blood pressure [81]. On the other hand, HIF stabilizers may raise EPO to more physiologic levels, reduce cross talk with vascular EPO receptors and mitigate hypertension. A 2014 study demonstrated that compared with EPO, treatment with the HIF stabilizer molidustat corrected anemia associated with subtotal nephrectomy, but in contrast to EPO, it reduced systolic BP in a dose-dependent manner [82]. The authors postulate that anti-inflammatory and antifibrotic effects of the drug on the kidney may be operative. This study further illustrates that increases in hematocrit can be dissociated from an increase in blood pressure.

Several HIF stabilizers are in Phase 3 clinical trials for the treatment of anemia in CKD. The ongoing Phase 3 trials of HIF stabilizers are evaluating, among patients with CKD, the equivalence of hemoglobin increase from baseline compared with

approved EPO-stimulating agents. Whereas hypertension is not a primary outcome measure for these Phase 3 trials, the equivalence of the cardiovascular outcomes is an important safety endpoint.

Human data are available but are inadequate to address the question of hypertension with HIF stabilizers. Among 145 patients with CKD not on dialysis who were treated with roxadustat for 16–24 weeks in varying doses, hypertension was reported in 11 (7.6%) [83]. Among 60 incident dialysis patients treated with roxadustat for 12 weeks, hypertension requiring an escalation of antihypertensive therapy occurred in 6 (10%) [84]. In the above studies, there was no placebo group or a group treated with EPO alone, therefore, whether the drug is equivalent to, safer or more detrimental than EPO or placebo with respect to hypertension remains unclear. In a 2017 study in China, among 61 CKD patients treated with roxadustat over 8 weeks, 4 (7%) developed hypertension compared with none of 30 treated with placebo [85]. In the same report, among 74 patients on dialysis treated for 6 weeks, 3 (4%) developed hypertension compared with 1 of 22 (5%) treated with EPO [85]. In a 2016 study of vadadustat among nondialysis CKD patients treated for 20 weeks, 138 were assigned to vadadustat and 72 to placebo; 11 in the vadadustat group (8%) and 2 in the placebo group (2.8%) had hypertension reported as an adverse event.

CONCLUSIONS

In summary, mediated by a variety of molecules, there is an imbalance in the vascular tone favoring net vasoconstriction that mediates hypertension due to EPO, especially in the setting of CKD. Besides the direct effects of the prostanoids, endothelins and NO pathways, EPO administration is also associated with increased responsiveness to catecholamines and angiotensin II and mitigation of hypoxia-induced vasodilation responses. EPO-induced hypertension, at least in part, appears to be independent of an increase in hemoglobin, because experiments show that hemoglobin may be increased by EPO without an increase in BP by simply treating the animals with EPO binding protein and that treatment with EPO in the setting of iron deficiency may not increase hemoglobin but may still increase BP. However, experimental data are not consistent across studies and better mechanistic designs are needed, especially in people with CKD, to dissect the precise mechanism of EPO-induced hypertension. Animal studies suggest that HIF stabilizers may provoke hypertension and those with high phosphorus concentrations and sleep apnea may be at an increased risk. Others show that there may be an antihypertensive effect via kidney repair. Whether this class of drugs will reduce the risk of hypertension compared with EPO remains to be seen.

CONFLICT OF INTEREST STATEMENT

R.A. serves on steering committees of clinical trials funded by Akebia, Johnson & Johnson, Relypsa, Bayer and Boehringer Ingelheim.

REFERENCES

1. Abraham PA, Macres MG. Blood pressure in hemodialysis patients during amelioration of anemia with erythropoietin. *J Am Soc Nephrol* 1991; 2: 927–936
2. van de Borne P, Tielemans C, Vanherweghem J-L *et al.* Effect of recombinant human erythropoietin therapy on ambulatory blood pressure and heart rate in chronic hemodialysis patients. *Nephrol Dial Transplant* 1992; 7: 45–49
3. Effect of recombinant human erythropoietin therapy on blood pressure in hemodialysis patients. Canadian Erythropoietin Study Group. *Am J Nephrol* 1991; 11: 23–26
4. Eschbach JW, Aquilino T, Haley NR *et al.* The long-term effects of recombinant human erythropoietin on the cardiovascular system. *Clin Nephrol* 1992; 38: S98–S103
5. Lebel M, Kingma I, Grose JH *et al.* Effect of recombinant human erythropoietin therapy on ambulatory blood pressure in normotensive and in untreated borderline hypertensive hemodialysis patients. *Am J Hypertens* 1995; 8: 545–551
6. Krapf R, Hulter HN. Arterial hypertension induced by erythropoietin and erythropoiesis-stimulating agents (ESA). *Clin J Am Soc Nephrol* 2009; 4: 470–480
7. Berns JS, Rudnick MR, Cohen RM *et al.* Effects of normal hematocrit on ambulatory blood pressure in epoetin-treated hemodialysis patients with cardiac disease. *Kidney Int* 1999; 56: 253–260
8. Conlon PJ, Kovalik E, Schumm D *et al.* Normalization of hematocrit in hemodialysis patients with cardiac disease does not increase blood pressure. *Ren Fail* 2000; 22: 435–444
9. Salem MM. Hypertension in the hemodialysis population: a survey of 649 patients. *Am J Kidney Dis* 1995; 26: 461–468
10. Poux JM, Lartigue M, Chaisemartin RA *et al.* Uraemia is necessary for erythropoietin-induced hypertension in rats. *Clin Exp Pharmacol Physiol* 1995; 22: 769–771
11. Lacasse M, Kingma I, Lariviere R *et al.* Uremia enhances the blood pressure response to erythropoietin. *Clin Exp Hypertens* 1997; 19: 389–401
12. Garcia DL, Anderson S, Rennke HG *et al.* Anemia lessens and its prevention with recombinant human erythropoietin worsens glomerular injury and hypertension in rats with reduced renal mass. *Proc Natl Acad Sci USA* 1988; 85: 6142–6146
13. Vaziri ND, Zhou XJ, Naqvi F *et al.* Role of nitric oxide resistance in erythropoietin-induced hypertension in rats with chronic renal failure. *Am J Physiol* 1996; 271: E113–E122
14. Kaupke CJ, Kim S, Vaziri ND. Effect of erythrocyte mass on arterial blood pressure in dialysis patients receiving maintenance erythropoietin therapy. *J Am Soc Nephrol* 1994; 4: 1874–1878
15. Buckner FS, Eschbach JW, Haley NR *et al.* Hypertension following erythropoietin therapy in anemic hemodialysis patients. *Am J Hypertens* 1990; 3: 947–955
16. Suzuki N, Ohneda O, Takahashi S *et al.* Erythroid-specific expression of the erythropoietin receptor rescued its null mutant mice from lethality. *Blood* 2002; 100: 2279–2288
17. Anagnostou A, Liu Z, Steiner M *et al.* Erythropoietin receptor mRNA expression in human endothelial cells. *Proc Natl Acad Sci USA* 1994; 91: 3974–3978
18. Carlini RG, Dusso AS, Obialo CI *et al.* Recombinant human erythropoietin (rHuEPO) increases endothelin-1 release by endothelial cells. *Kidney Int* 1993; 43: 1010–1014
19. Anagnostou A, Lee ES, Kessimian N *et al.* Erythropoietin has a mitogenic and positive chemotactic effect on endothelial cells. *Proc Natl Acad Sci USA* 1990; 87: 5978–5982
20. Nielsen OJ. Pharmacokinetics of recombinant human erythropoietin in chronic haemodialysis patients. *Pharmacol Toxicol* 1990; 66: 83–86
21. Kagaya Y, Asaumi Y, Wang W *et al.* Current perspectives on protective roles of erythropoietin in cardiovascular system: erythropoietin receptor as a novel therapeutic target. *Tohoku J Exp Med* 2012; 227: 83–91
22. Tada H, Kagaya Y, Takeda M *et al.* Endogenous erythropoietin system in non-hematopoietic lineage cells plays a protective role in myocardial ischemia/reperfusion. *Cardiovasc Res* 2006; 71: 466–477

23. Asaumi Y, Kagaya Y, Takeda M *et al.* Protective role of endogenous erythropoietin system in nonhematopoietic cells against pressure overload-induced left ventricular dysfunction in mice. *Circulation* 2007; 115: 2022–2032
24. Satoh K, Kagaya Y, Nakano M *et al.* Important role of endogenous erythropoietin system in recruitment of endothelial progenitor cells in hypoxia-induced pulmonary hypertension in mice. *Circulation* 2006; 113: 1442–1450
25. Nakano M, Satoh K, Fukumoto Y *et al.* Important role of erythropoietin receptor to promote VEGF expression and angiogenesis in peripheral ischemia in mice. *Circ Res* 2007; 100: 662–669
26. Lee MS, Lee JS, Lee JY. Prevention of erythropoietin-associated hypertension. *Hypertension* 2007; 50: 439–445
27. Burrill DR, Vernet A, Collins JJ *et al.* Targeted erythropoietin selectively stimulates red blood cell expansion in vivo. *Proc Natl Acad Sci USA* 2016; 113: 5245–5250
28. Schmieder RE, Langenfeld MR, Hilgers KF. Endogenous erythropoietin correlates with blood pressure in essential hypertension. *Am J Kidney Dis* 1997; 29: 376–382
29. Heidenreich S, Rahn KH, Zidek W. Direct vasopressor effect of recombinant human erythropoietin on renal resistance vessels. *Kidney Int* 1991; 39: 259–265
30. Neusser M, Tepel M, Zidek W. Erythropoietin increases cytosolic free calcium concentration in vascular smooth muscle cells. *Cardiovasc Res* 1993; 27: 1233–1236
31. Hon G, Vaziri ND, Kaupke CJ *et al.* Lack of a fast-acting effect of erythropoietin on arterial blood pressure and endothelin level. *Artif Organs* 1995; 19: 188–191
32. Miyashita K, Tojo A, Kimura K *et al.* Blood pressure response to erythropoietin injection in hemodialysis and predialysis patients. *Hypertens Res* 2004; 27: 79–84
33. Briet M, Barhoumi T, Mian MO *et al.* Effects of recombinant human erythropoietin on resistance artery endothelial function in stage 4 chronic kidney disease. *J Am Heart Assoc* 2013; 2: e000128
34. D'Amours M, Chbinou N, Beaudoin J *et al.* Increased ET-1 and reduced ET(B) receptor expression in uremic hypertensive rats. *Clin Exp Hypertens* 2010; 32: 61–69
35. Lariviere R, Lebel M. Endothelin-1 in chronic renal failure and hypertension. *Can J Physiol Pharmacol* 2003; 81: 607–621
36. Lebel M, Rodrigue ME, Agharazii M *et al.* Antihypertensive and renal protective effects of renin-angiotensin system blockade in uremic rats treated with erythropoietin. *Am J Hypertens* 2006; 19: 1286–1292
37. Bode-Boger SM, Boger RH, Kuhn M *et al.* Recombinant human erythropoietin enhances vasoconstrictor tone via endothelin-1 and constrictor prostanoids. *Kidney Int* 1996; 50: 1255–1261
38. Zhou XJ, Pandian D, Wang XQ *et al.* Erythropoietin-induced hypertension in rat is not mediated by alterations of plasma endothelin, vasopressin, or atrial natriuretic peptide levels. *J Am Soc Nephrol* 1997; 8: 901–905
39. Lebel M, Lacasse M, Lariviere R *et al.* Plasma and blood vessel endothelin-1 concentrations in hypertensive uremic rats treated with erythropoietin. *Clin Exp Hypertens* 1998; 20: 939–951
40. Moreau C, Lariviere R, Kingma I *et al.* Chronic nitric oxide inhibition aggravates hypertension in erythropoietin-treated renal failure rats. *Clin Exp Hypertens* 2000; 22: 663–674
41. Rancourt ME, Rodrigue ME, Agharazii M *et al.* Role of oxidative stress in erythropoietin-induced hypertension in uremic rats. *Am J Hypertens* 2010; 23: 314–320
42. Barhoumi T, Briet M, Kasal DA *et al.* Erythropoietin-induced hypertension and vascular injury in mice overexpressing human endothelin-1: exercise attenuated hypertension, oxidative stress, inflammation and immune response. *J Hypertens* 2014; 32: 784–794
43. Rodrigue ME, Brochu E, Orleans-Juste P *et al.* Effect of erythropoietin on blood pressure and on the vascular endothelial ET-1/ETB receptor system. *Am J Hypertens* 2008; 21: 639–643
44. Brochu E, Lacasse S, Lariviere R *et al.* Differential effects of endothelin-1 antagonists on erythropoietin-induced hypertension in renal failure. *J Am Soc Nephrol* 1999; 10: 1440–1446
45. Muller R, Steffen HM, Brunner R *et al.* Changes in the alpha adrenergic system and increase in blood pressure with recombinant human erythropoietin (rHuEpo) therapy for renal anemia. *Clin Invest Med* 1991; 14: 614–622
46. Takahashi K, Totsune K, Imai Y *et al.* Plasma concentrations of immunoreactive-endothelin in patients with chronic renal failure treated with recombinant human erythropoietin. *Clin Sci* 1993; 84: 47–50
47. Stefanidis I, Mertens PR, Wurth P *et al.* Influence of recombinant human erythropoietin therapy on plasma endothelin-1 levels during hemodialysis. *Int J Artif Organs* 2001; 24: 367–373
48. Kang DH, Yoon KI, Han DS. Acute effects of recombinant human erythropoietin on plasma levels of proendothelin-1 and endothelin-1 in haemodialysis patients. *Nephrol Dial Transplant* 1998; 13: 2877–2883
49. Lebel M, Grose JH, Kingma I *et al.* Plasma endothelin levels and blood pressure in hemodialysis and in CAPD patients. Effect of subcutaneous erythropoietin replacement therapy. *Clin Exp Hypertens* 1994; 16: 565–575
50. Lebel M, Moreau V, Grose JH *et al.* Plasma and peritoneal endothelin levels and blood pressure in CAPD patients with or without erythropoietin replacement therapy. *Clin Nephrol* 1998; 49: 313–318
51. Yamakado M, Umezu M, Nagano M *et al.* Mechanisms of hypertension induced by erythropoietin in patients on hemodialysis. *Clin Invest Med* 1991; 14: 623–629
52. Rodrigue ME, Lacasse M, Lariviere R *et al.* Cyclooxygenase inhibition with acetylsalicylic acid unmasks a role for prostacyclin in erythropoietin-induced hypertension in uremic rats. *Can J Physiol Pharmacol* 2005; 83: 467–475
53. Bode-Boger SM, Boger RH, Kuhn M *et al.* Endothelin release and shift in prostaglandin balance are involved in the modulation of vascular tone by recombinant erythropoietin. *J Cardiovasc Pharmacol* 1992; 20: S25–S28
54. Rodrigue ME, Moreau C, Lariviere R *et al.* Relationship between eicosanoids and endothelin-1 in the pathogenesis of erythropoietin-induced hypertension in uremic rats. *J Cardiovasc Pharmacol* 2003; 41: 388–395
55. Caravaca F, Pizarro JL, Arrobas M *et al.* Antiplatelet therapy and development of hypertension induced by recombinant human erythropoietin in uremic patients. *Kidney Int* 1994; 45: 845–851
56. Wang XQ, Vaziri ND. Erythropoietin depresses nitric oxide synthase expression by human endothelial cells. *Hypertension* 1999; 33: 894–899
57. Scalera F, Kielstein JT, Martens-Lobenhoffer J *et al.* Erythropoietin increases asymmetric dimethylarginine in endothelial cells: role of dimethylarginine dimethylaminohydrolase. *J Am Soc Nephrol* 2005; 16: 892–898
58. Teerlink T. ADMA metabolism and clearance. *Vasc Med* 2005; 10 (Suppl 1): S73–S81
59. Desai A, Zhao Y, Warren JS. Human recombinant erythropoietin augments serum asymmetric dimethylarginine concentrations but does not compromise nitric oxide generation in mice. *Nephrol Dial Transplant* 2008; 23: 1513–1520
60. Vaziri ND, Zhou XJ, Naqvi F *et al.* Role of nitric oxide resistance in erythropoietin-induced hypertension in rats with chronic renal failure. *Am J Physiol* 1996; 271: E113–E122
61. Wilcox CS, Deng X, Doll AH *et al.* Nitric oxide mediates renal vasodilation during erythropoietin-induced polycythemia. *Kidney Int* 1993; 44: 430–435
62. del CD, Raji L, Shultz PJ *et al.* The pressor effect of recombinant human erythropoietin is not due to decreased activity of the endogenous nitric oxide system. *Nephrol Dial Transplant* 1995; 10: 505–508
63. Ni Z, Wang XQ, Vaziri ND. Nitric oxide metabolism in erythropoietin-induced hypertension: effect of calcium channel blockade. *Hypertension* 1998; 32: 724–729
64. Tsukahara H, Hiraoka M, Hori C *et al.* Chronic erythropoietin treatment enhances endogenous nitric oxide production in rats. *Scand J Clin Lab Invest* 1997; 57: 487–493
65. Migliori M, Taccola D, Panichi V *et al.* Nitric oxide-dependent renal vasodilation is not altered in rat with rHuEpo-induced hypertension. *Kidney Blood Press Res* 1999; 22: 140–145
66. Savard S, Lavoie P, Villeneuve C *et al.* eNOS gene delivery prevents hypertension and reduces renal failure and injury in rats with reduced renal mass. *Nephrol Dial Transplant* 2012; 27: 2182–2190
67. Ruschitzka FT, Wenger RH, Stallmach T *et al.* Nitric oxide prevents cardiovascular disease and determines survival in polyglobulic mice overexpressing erythropoietin. *Proc Natl Acad Sci USA* 2000; 97: 11609–11613
68. Ioka T, Tsuruoka S, Ito C *et al.* Hypertension induced by erythropoietin has a correlation with truncated erythropoietin receptor mRNA in endothelial

- progenitor cells of hemodialysis patients. *Clin Pharmacol Ther* 2009; 86: 154–159
69. Ksiazek A, Zaluska WT, Ksiazek P. Effect of recombinant human erythropoietin on adrenergic activity in normotensive hemodialysis patients. *Clin Nephrol* 2001; 56: 104–110
 70. Hori K, Onoyama K, Iseki K *et al*. Hemodynamic and volume changes by recombinant human erythropoietin (rHuEPO) in the treatment of anemic hemodialysis patients. *Clin Nephrol* 1990; 33: 293–298
 71. Hand MF, Haynes WG, Johnstone HA *et al*. Erythropoietin enhances vascular responsiveness to norepinephrine in renal failure. *Kidney Int* 1995; 48: 806–813
 72. Muller R, Steffen HM, Brunner R *et al*. [Disordered alpha 2-adrenoreceptor function in hemodialysis patients with renal anemia—a possible cause of increased blood pressure in relation to recombinant human erythropoietin?]. *Klin Wochenschr* 1991; 69: 742–748
 73. Eggena P, Willsey P, Jamgotchian N *et al*. Influence of recombinant human erythropoietin on blood pressure and tissue renin-angiotensin systems. *Am J Physiol* 1991; 261: E642–E646
 74. Barrett JD, Zhang Z, Zhu JH *et al*. Erythropoietin upregulates angiotensin receptors in cultured rat vascular smooth muscle cells. *J Hypertens* 1998; 16: 1749–1757
 75. Brier ME, Bunke CM, Lathon PV *et al*. Erythropoietin-induced antinatriuresis mediated by angiotensin II in perfused kidneys. *J Am Soc Nephrol* 1993; 3: 1583–1590
 76. Lundby C, Thomsen JJ, Boushel R *et al*. Erythropoietin treatment elevates haemoglobin concentration by increasing red cell volume and depressing plasma volume. *J Physiol* 2007; 578: 309–314
 77. Lebel M, Kingma I, Grose JH *et al*. Hemodynamic and hormonal changes during erythropoietin therapy in hemodialysis patients. *J Am Soc Nephrol* 1998; 9: 97–104
 78. Roger SD, Grasty MS, Baker LR *et al*. Effects of oxygen breathing and erythropoietin on hypoxic vasodilation in uremic anemia. *Kidney Int* 1992; 42: 975–980
 79. Semenza GL. Hypoxia-inducible factors in physiology and medicine. *Cell* 2012; 148: 399–408
 80. Mokas S, Lariviere R, Lamallice L *et al*. Hypoxia-inducible factor-1 plays a role in phosphate-induced vascular smooth muscle cell calcification. *Kidney Int* 2016; 90: 598–609
 81. Agarwal R, Light RP. Arterial stiffness and interdialytic weight gain influence ambulatory blood pressure patterns in hemodialysis patients. *Am J Physiol Renal Physiol* 2007; 294: F303–F308
 82. Flamme I, Oehme F, Ellinghaus P *et al*. Mimicking hypoxia to treat anemia: HIF-stabilizer BAY 85-3934 (Molidustat) stimulates erythropoietin production without hypertensive effects. *PLoS One* 2014; 9: e111838
 83. Provenzano R, Besarab A, Sun CH *et al*. Oral hypoxia-inducible factor prolyl hydroxylase inhibitor roxadustat (FG-4592) for the treatment of anemia in patients with CKD. *Clin J Am Soc Nephrol* 2016; 11: 982–991
 84. Besarab A, Chernyavskaya E, Motylev I *et al*. Roxadustat (FG-4592): correction of anemia in incident dialysis patients. *J Am Soc Nephrol* 2016; 27: 1225–1233
 85. Chen N, Qian J, Chen J *et al*. Phase 2 studies of oral hypoxia-inducible factor prolyl hydroxylase inhibitor FG-4592 for treatment of anemia in China. *Nephrol Dial Transplant* 2017; 32: 1373–1386

Received: 9.8.2017; Editorial decision: 23.10.2017